

this product if you are taking sedatives or tranquilizers, without first consulting your doctor. Use caution when driving a motor vehicle or operating machinery.”

(8) *For products containing diphenhydramine hydrochloride identified in § 336.10(c).* “Do not use [bullet]¹ with any other product containing diphenhydramine, including one used on skin”.

(d) *Directions.* The labeling of the product contains the following information under the heading “Directions”:

(1) *For products containing cyclizine hydrochloride identified in § 336.10(a).* Adults and children 12 years of age and over: Oral dosage is 50 milligrams every 4 to 6 hours, not to exceed 200 milligrams in 24 hours, or as directed by a doctor. Children 6 to under 12 years of age: Oral dosage is 25 milligrams every 6 to 8 hours, not to exceed 75 milligrams in 24 hours, or as directed by a doctor.

(2) *For products containing dimenhydrinate identified in § 336.10(b).* Adults and children 12 years of age and over: Oral dosage is 50 to 100 milligrams every 4 to 6 hours, not to exceed 400 milligrams in 24 hours, or as directed by a doctor. Children 6 to under 12 years of age: Oral dosage is 25 to 50 milligrams every 6 to 8 hours, not to exceed 150 milligrams in 24 hours, or as directed by a doctor. Children 2 to under 6 years of age: Oral dosage is 12.5 to 25 milligrams every 6 to 8 hours, not to exceed 75 milligrams in 24 hours, or as directed by a doctor.

(3) *For products containing diphenhydramine hydrochloride identified in § 336.10(c).* Adults and children 12 years of age and over: Oral dosage is 25 to 50 milligrams every 4 to 6 hours, not to exceed 300 milligrams in 24 hours, or as directed by a doctor. Children 6 to under 12 years of age: Oral dosage is 12.5 to 25 milligrams every 4 to 6 hours, not to exceed 150 milligrams in 24 hours, or as directed by a doctor.

(4) *For products containing meclizine hydrochloride identified in § 336.10(d).* Adults and children 12 years of age and

over: Oral dosage is 25 to 50 milligrams once daily, or as directed by a doctor.

(e) The word “physician” may be substituted for the word “doctor” in any of the labeling statements in this section.

[52 FR 15892, Apr. 30, 1987, as amended at 53 FR 35809, Sept. 15, 1988; 59 FR 16982, Apr. 11, 1994; 67 FR 72559, Dec. 6, 2003]

§ 336.80 Professional labeling.

The labeling provided to health professionals (but not to the general public) may contain the following additional indications.

(a) *For products containing cyclizine hydrochloride, dimenhydrinate, and diphenhydramine hydrochloride identified in § 336.10 (a), (b), and (c).* “For the treatment of vertigo of motion sickness.”

(b) *For products containing meclizine hydrochloride identified in § 336.10(d).* “For the treatment of vertigo.”

PART 338—NIGHTTIME SLEEP-AID DRUG PRODUCTS FOR OVER-THE-COUNTER HUMAN USE

Subpart A—General Provisions

Sec.
338.1 Scope.
338.3 Definition.

Subpart B—Active Ingredients

338.10 Nighttime sleep-aid active ingredients.

Subpart C—Labeling

338.50 Labeling of nighttime sleep-aid drug products.

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371.

SOURCE: 54 FR 6826, Feb. 14, 1989, unless otherwise noted.

Subpart A—General Provisions

§ 338.1 Scope.

(a) An over-the-counter nighttime sleep-aid drug product in a form suitable for oral administration is generally recognized as safe and effective and is not misbranded if it meets each condition in this part and each general condition established in § 330.1 of this chapter.

¹ See § 201.66(b)(4) of this chapter for definition of bullet symbol.